

AD-A080 737 VIRGINIA POLYTECHNIC INST AND STATE UNIV BLACKSBURG --ETC F/8 12/1
SOME RESULTS ON SOJOURN TIMES IN ACYCLIC JACKSON NETWORKS. REVI--ETC(U)
DEC 78 B SIMON, R D FOLEY N00014-77-C-0743
UNCLASSIFIED VTR-79-01-REV NL

1 x 1
AD-A080737



END
DATE
FILMED
3 - 80
100

ADA 080737

LIVEL (R)
SOME RESULTS ON SOJOURN TIMES
IN ACYCLIC JACKSON NETWORKS

by

B. Simon

R. D. Foley

SOME RESULTS ON SOJOURN TIMES
IN ACYCLIC JACKSON NETWORKS

by


B. Simon

R. D. Foley

Department of Industrial Engineering and Operations Research
Virginia Polytechnic Institute and State University
Blacksburg, Virginia

TECH REPORT
VTR - 79-01
July, 1978
Revised, December 1978.

This research was supported jointly by
NSF Grant ENG77-22757 and by the
Office of Naval Research Contract
N00014-77-C-0743 (NR042-296)



Distribution of this document is unlimited. Reproduction in whole or
in part is permitted for any purpose of the United State Government.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
9 <u>Technical Report VTR 79-01</u>		
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED	
6 <u>SOME RESULTS ON SOJOURN TIMES IN ACYCLIC JACKSON NETWORKS - REVISION</u>	<u>Technical Report 1979</u>	
7. AUTHOR(s)	14 <u>VTR-79-01-REV</u>	
10 <u>Burton/Simon ^{D.} Robert/Foley</u>	8. CONTRACT OR GRANT NUMBER(s)	
9. PERFORMING ORGANIZATION NAME AND ADDRESS	15 <u>NO0014-77-C-0743</u>	
<u>Dept. of Industrial Engineering & Operations Res. Virginia Polytechnic Inst. & State University Blacksburg, Virginia 24061</u>	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <u>NSF-ENG 77-22757</u>	
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE	
<u>Director, Statistics and Probability Program Mathematical & Information Sciences Division 800 N. Quincy St., Arlington, VA 22217</u>	<u>December 1979</u>	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES	
	<u>14</u>	
	15. SECURITY CLASS. (of this report)	
	<u>Unclassified</u>	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)		
<u>Approved for Public Release; Distribution Unlimited</u>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
<u>11/ Dec 78</u>		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
<u>Jackson Networks Sojourn Times</u>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
<u>Lemoine (1977) studies sojourn times in acyclic, Jackson networks as part of his survey of equilibrium results. In his paper, he argues that a particular customer has a sojourn time at node i that is independent of his remaining sojourn time in the network given that the customer transfer from node i to node j. This assumption is then used to derive a set of equations involving the Laplace-Stieltjes transform of the sojourn times at each node from which several properties of sojourn times are determined.</u>		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

406 747

JM

Abstract

~~Lemoine (1977)~~ studies ^λsojourn times in acyclic, Jackson networks as part of ^λhis survey of equilibrium results. ^{Argument is this} ~~In his paper, he argues~~ that a particular customer has a sojourn time at node i that is independent of his remaining sojourn time in the network given that the customer transfer from node i to node j. This assumption is then used to derive a set of equations involving the Laplace-Stieltjes transform of the sojourn times at each node from which several properties of sojourn times are determined. ~~✓~~

In a private communication, Mitrani (1978) argues that Lemoine's independence assumption is in error and provides a heuristic argument based on a four node, acyclic, Jackson network. From this, Mitrani concludes that the sojourn time in such networks is still an unsolved problem. We make Mitrani's argument rigorous, using some new results on sojourn times in Jackson networks.

Accession For	
NY 100-100000	<input checked="" type="checkbox"/>
Dis. Ill.	<input type="checkbox"/>
Unsub. and	<input type="checkbox"/>
Sub. and Ill.	<input type="checkbox"/>
For Dis. Ill. Sub. and Ill. Codes Sub. and/or Special	

Continuation of Abstract.

In a private communication, Mitrani (1978) argues that Lemoine's independence assumption is in error and provides a heuristic argument based on a four node, acyclic, Jackson network. From this, Mitrani concludes that the sojourn time in such networks is still an unsolved problem. We make Mitrani's argument rigorous, using some new results on sojourn times in Jackson networks.

SOME RESULTS ON SOJOURN TIMES

IN ACYCLIC JACKSON NETWORKS

1. Background. Lemoine (1977) analyzes the sojourn time of a customer in an acyclic Jackson network. His analysis relies on the sojourn time of a particular customer at server i being independent of the customer's remaining sojourn time in the network, given the customer transfers from server i to server j . Using this independence assumption, Lemoine derives the Laplace-Stieltjes transform for the sojourn time of an arbitrary customer in the network.

In a private communication, Mitrani (1978) challenges Lemoine's assertion that the sojourn time at server i and the remaining sojourn time in the network are independent given that the customer transfers from i to j . In particular he concludes that the distribution of the sojourn time of the customer in an acyclic Jackson network is still an unsolved problem.

We resolve these issues in this paper. Through the use of a counter-example, we substantiate Mitrani's claim. This work culminates with Theorem 4.3. In section 5 we will summarize our results very briefly and point out related literature. We assume the reader is familiar with Jackson (1957) and Lemoine (1977).

2. Statement of Problem. In this section we construct a three node, acyclic Jackson network with the property that a customer's remaining sojourn time is not independent of his sojourn time in node 1, given that he goes through node 2. The network has the following form.




Figure 2.1

There is a Poisson stream with rate λ arriving at node 1. When a customer leaves node 1 he goes directly to node 3 with probability $(1-p)$, or goes to node 2 with probability p . If the customer goes to node 2, he goes directly to node 3 when his service is completed. The customer leaves the network after service at node 3. Service times at node 1 are exponentially distributed with parameter μ_1 .

To show that Lemoine's argument in section 1 is faulty, it suffices to show that the sojourn time in the first queue, T_1 , and the total sojourn time in the second and third queue, $T_2 + T_3$, are dependent random variables for a customer who goes through the second queue. For such a customer we will show that for certain values of λ , μ_1 , μ_2 , μ_3 , p , and t ; the expected sojourn time in the third queue given that the sojourn time in the first queue is t , $E[T_3|T_1=t]$, is greater than the unconditional expected sojourn time in the third queue $E[T_3]$. By Theorem 3.3, below, the expected sojourn time in the second queue is independent of T_1 . Thus, for the above parameters,

$$\begin{aligned} E(T_3 + T_2 | T_1 = t) &= E(T_3 | T_1 = t) + E(T_2) \\ &> E(T_3) + E(T_2) = E(T_3 + T_2). \end{aligned} \tag{1}$$

This fact verifies that the remaining sojourn time (after leaving queue 1) and the sojourn time in queue 1 are not independent. In turn this verifies the Mitrani conjecture.

3. Preliminary Results. In order to formally show that the sojourn times in the first and third queues are dependent, we need the following results. Theorem 3.3 is proven in more generality than is necessary for the problem at hand, but gives an intuitive reason for believing that sojourn times in Jackson networks are not independent in general.

Let J be an acyclic Jackson network with nodes $(1, 2, \dots, N)$ and switching matrix P . Let

λ_i = exogenous arrival rate to node i ,

α_i = total arrival rate to node i (which is given by the solution of the traffic equation),

μ_i = service rate at node i .

$M_i(s)$ = the number of customers at node i at time s ,

For a fixed customer ("our customer") who goes through nodes (r_1, r_2, \dots, r_k) , let

a_i = the time of arrival at node i ,

t_i = the time of departure from node i ,

$T_i = t_i - a_i$ = the sojourn time of our customer at node i ,

$D_i(s)$ = the number of departures from node i in the interval $(t_i, t_i + s]$,

$D_i^*(s)$ = the number of departures from node i in the interval $(a_i, a_i + s]$.

Lemma 3.1 Let J be a steady state, acyclic, Jackson network. Suppose our customer enters node j from outside the system (an exogenous arrival). Then

$$P(D_j^*(s) = k | T_j = t) = \frac{(\alpha_j s)^k e^{-\alpha_j s}}{k!}, \quad \forall s \in (0, T_j).$$

Proof.

$$\begin{aligned} P(D_j^*(s) = k | T_j = t) &= \sum_{m=k}^{\infty} P(D_j^*(s) = k | T_j = t, M_j(a_j^-) = m) P(M_j(a_j^-) = m | T_j = t) \\ &= \sum_{m=k}^{\infty} \binom{m}{k} \left(\frac{s}{t}\right)^k \left(\frac{t-s}{t}\right)^{m-k} \frac{\mu_j (\mu_j t)^m e^{-\mu_j t}}{m!} \frac{\left(1 - \frac{\alpha_j}{\mu_j}\right) \left(\frac{\alpha_j}{\mu_j}\right)^m}{(\mu_j - \alpha_j) e^{-(\mu_j - \alpha_j)t}} \\ &= \frac{e^{-\alpha_j t}}{k!} \left(\frac{s}{t-s}\right)^k \sum_{m=k}^{\infty} \frac{(\alpha_j (t-s))^m}{(m-k)!} \\ &= \frac{(\alpha_j s)^k e^{-\alpha_j s}}{k!} \cdot \square \end{aligned}$$

Using Theorem 1.18 Çinlar (1975:p.77), one immediately has

Corollary 3.2. The departure process from node j is a Poisson process with rate α_j in (a_j, t_j) for any T_j .

This result also follows from the Sojourn Time Theorem in Burke (1972).

Theorem 3.3. Suppose an exogenous arrival enters node j (of a steady state, acyclic, Jackson network) at time a_j . Then

$$\begin{aligned} P(M_1(t_j^-) = k_1, M_2(t_j^-) = k_2, \dots, M_j(t_j^-) = k_j + 1, \dots, M_N(t_j^-) = k_N | T_j = t) \\ = \frac{(\alpha_j t)^{k_j} e^{-\alpha_j t}}{k_j!} \prod_{i \neq j} \left(1 - \frac{\alpha_i}{\mu_i}\right) \left(\frac{\alpha_i}{\mu_i}\right)^{k_i} \end{aligned} \quad (2)$$

Proof. Let J_2 be the set of nodes which can be reached after leaving node j either directly or after passing through other nodes in J_2 . Let $J_1 = (\{j\} \cup J_2)^c$. Without loss of generality, assume the nodes in J_1 are numbered $1, \dots, j-1$ and the nodes in J_2 are $j+1, \dots, N$. We will call a customer flow a departure process from J_1 if it originates at a node $k \in J_1$ and either leaves the system entirely or goes to some node $m \notin J_1$.

At time a_j^- the network is in steady state. Hence,

$$P\{M_1(a_j^-) = k_1, \dots, M_N(a_j^-) = k_N\} = \prod_{i=1}^N \left(1 - \frac{\alpha_i}{\mu_i}\right) \left(\frac{\alpha_i}{\mu_i}\right)^{k_i}$$

The arrival processes to J_1 after time a_j are Poisson processes independent of T_j . In addition, the service times of customers in J_1 after time a_j are independent of T_j .

Thus, $\forall s \in (a_j, \infty)$

$$P\{M_1(s) = k_1, \dots, M_{j-1}(s) = k_{j-1} | T_j = t\} = \prod_{i=1}^{j-1} \left(1 - \frac{\alpha_i}{\mu_i}\right) \left(\frac{\alpha_i}{\mu_i}\right)^{k_i}.$$

In particular for $s = t_j^-$, we have

$$P\{M_1(t_j^-) = k_1, \dots, M_{j-1}(t_j^-) = k_{j-1} | T_j = t\} = \prod_{i=1}^{j-1} \left(1 - \frac{\alpha_i}{\mu_i}\right) \left(\frac{\alpha_i}{\mu_i}\right)^{k_i}.$$

Since the network is acyclic, by Beutler and Melamed (1978) the departure processes from J_1 before time t_j are mutually independent Poisson processes, and independent of the state of the queues in J_1 at time t_j^- . Hence,

$$P\{M_j(t_j^-) = k_j + 1 | T_j = t, M_1(t_j^-) = k_1, \dots, M_{j-1}(t_j^-) = k_{j-1}\} = \frac{e^{-\alpha_j t} (\alpha_j t)^{k_j}}{k_j!}.$$

Let $D_j'(s)$ be the number of departures from node j in the interval $(a_j, a_j + s]$. For $s < T_j$, $D_j'(s)$ is completely determined by $M_j(a_j^-)$, T_j and the service times at j . Since $M_j(a_j^-)$, T_j and the service times at queue j are independent of the departure processes from J_1 in the interval (a_j, t_j^-) , $D_j'(s)$ is independent of the departure processes from J_1 during (a_j, t_j^-) . From Corollary 3.2, $D_j'(s)$ is a Poisson process. Thus $D_j'(s)$ and the departure processes from J_1 before time t_j are mutually independent Poisson processes. Thus the arrival processes to J_2 are independent of T_j and we conclude that:

$$\begin{aligned} P\{M_{j+1}(t_j^-) = k_{j+1}, \dots, M_N(t_j^-) = k_N | T_j = t, M_1(t_j^-) = k_1, \dots, M_j(t_j^-) = k_j + 1\} \\ = \prod_{i=j+1}^N \left(1 - \frac{\alpha_i}{\mu_i}\right) \left(\frac{\alpha_i}{\mu_i}\right)^{k_i}. \end{aligned}$$

Combining these conditional probabilities, yields our result. \square

Since an exogenous arrival sees a steady state queue length distribution,

$$\text{it is well known that } P(T_j \leq t) = 1 - e^{-(\mu_j - \alpha_j)t}. \quad (3)$$

By unconditioning (2) by (3) over all possible values of t , one gets the usual Jackson steady state results.

4. Main Results. Our main result is the construction of an acyclic Jackson network where a customer's sojourn times at the queues in his path are not independent random variables. It should be clear from the construction that acyclic networks with independent sojourn times are the exception and not the rule.

Consider the network described in 2. Choose $\lambda > 0$, $p \in (0,1)$, $\mu_2 > 0$, $\mu_3 > 0$ so that $p\lambda < \mu_2$ and $\lambda < \mu_3$. From Disney, et al. (1973), the arrival stream to node 2 is Poisson with rate $p\lambda$ and the stream of customers going directly from node 1 to node 3 is Poisson with rate $(1-p)\lambda$. Thus $E(T_3)$ does not depend on the value of μ_1 . We will show that with a suitable choice of μ_1 and t we can make $E(T_3|T_1 = t)$ arbitrarily large. In particular we can make $E(T_3|T_1 = t) > E(T_3)$ so T_1 and T_3 are not independent.

Fix $r > 0$ and choose

$$\mu_1 > \text{Max} \left\{ \lambda, \left(r + \frac{1}{\mu_2 - p\lambda} \right) \frac{\mu_3 \mu_2^2}{(\mu_2 - p\lambda)(1 - e^{-\mu_2} - \mu_2 e^{-\mu_2})(1-p)} \right\}. \quad (4)$$

The values of λ , μ_1 , μ_2 , μ_3 and p assure that a steady state exists. Now fix $\epsilon > 0$ and choose n^* large enough so that

$$\sum_{j=0}^{n^*} \frac{(\mu_1 s)^j e^{-\mu_1 s}}{j!} > 1 - \epsilon, \quad \forall s \in (0,1]. \quad (5)$$

Since (5) is bounded below at $s = 1$, it suffices to choose n^* so that

$$\sum_{j=0}^{n^*} \frac{\mu_1^j e^{-\mu_1}}{j!} > 1 - \epsilon.$$

Given ϵ and n^* , now choose t large enough so that

$$P(M_1(t_1^+) > n^* | T_1 = t) > 1 - \epsilon. \quad (6)$$

This can be done since

$$P(M_1(t_1^+) > n^* | T_1 = t) = \sum_{j=n^*+1}^{\infty} \frac{(\lambda t)^j e^{-\lambda t}}{j!} \rightarrow 1 \text{ as } t \rightarrow \infty.$$

Let

D = number of departures from node 1 in $(t_1, a_3]$.

D^* = number of departures from node 3 in $(t_1, a_3]$.

Lemma 4.1 $E(D | T_1 = t) > (1 - \epsilon)^2 (\mu_2 - p\lambda) \frac{\mu_1}{\mu_2} (1 - e^{-\mu_2} - \mu_2 e^{-\mu_2}).$

Proof. $E(D | T_1 = t) = \sum_{j=0}^{\infty} E(D | T_1 = t, M_2(t_1^-) = j) P(M_2(t_1^-) = j | T_1 = t).$

By Theorem 3.3, $P(M_2(t_1^-) = j | T_1 = t) = \left(\frac{p\lambda}{\mu_2}\right)^j \left(1 - \frac{p\lambda}{\mu_2}\right)$, so

$$\begin{aligned} E(D | T_1 = t) &= \sum_{j=0}^{\infty} E(D | T_1 = t, M_2(t_1^-) = j) \left(\frac{p\lambda}{\mu_2}\right)^j \left(1 - \frac{p\lambda}{\mu_2}\right) \\ &> \left(1 - \frac{p\lambda}{\mu_2}\right) E(D | T_1 = t, M_2(t_1^-) = 0) = \frac{\mu_2 - p\lambda}{\mu_2} \int_0^{\infty} E(D_1(s) | T_1 = t) \mu_2 e^{-\mu_2 s} ds \\ &= (\mu_2 - p\lambda) \int_0^{\infty} \sum_{j=0}^{\infty} E(D_1(s) | T_1 = t, M_1(t_1^+) = j) P(M_1(t_1^+) = j | T_1 = t) e^{-\mu_2 s} ds. \end{aligned}$$

Since $D_1(s)$ is conditionally independent of T_1 given M_1 , we have

$$\begin{aligned} E(D|T_1 = t) &> (\mu_2 - p\lambda) \int_0^\infty \sum_{j=0}^\infty E(D_1(s) | M_1(t_1^+) = j) P(M_1(t_1^+) = j | T_1 = t) e^{-\mu_2 s} ds \\ &> (\mu_2 - p\lambda) \int_0^\infty \sum_{j=n^*+1}^\infty E(D_1(s) | M_1(t_1^+) = j) P(M_1(t_1^+) = j | T_1 = t) e^{-\mu_2 s} ds \end{aligned}$$

and since $E(D_1(s) | M_1(t_1^+) = j)$ is an increasing function of j ,

$$\geq (\mu_2 - p\lambda) \int_0^\infty \sum_{j=n^*+1}^\infty E(D_1(s) | M_1(t_1^+) = n^*) P(M_1(t_1^+) = j | T_1 = t) e^{-\mu_2 s} ds,$$

and by (6) this is

$$\begin{aligned} &> (1 - \epsilon) (\mu_2 - p\lambda) \int_0^\infty E(D_1(s) | M_1(t_1^+) = n^*) e^{-\mu_2 s} ds \\ &> (1 - \epsilon) (\mu_2 - p\lambda) \int_0^1 E(D_1(s) | M_1(t_1^+) = n^*) e^{-\mu_2 s} ds. \end{aligned}$$

Note that, $E(D_1(s) | M_1(t_1^+) = n^*) = \sum_{j=1}^\infty j P(D_1(s) = j | M_1(t_1^+) = n^*)$

$$\begin{aligned} &> \sum_{j=1}^{n^*} j P(D_1(s) = j | M_1(t_1^+) = n^*) \\ &= \sum_{j=1}^{n^*} \frac{(\mu_1 s)^j e^{-\mu_1 s}}{(j-1)!} \end{aligned}$$

which by (5) is $> (1 - \epsilon)\mu_1 s$. Thus,

$$\begin{aligned}
E(D|T_1 = t) &> (1 - \epsilon)^2 (\mu_2 - p\lambda) \int_0^1 \mu_1 s e^{-\mu_2 s} ds \\
&= (1 - \epsilon)^2 (\mu_2 - p\lambda) \frac{\mu_1}{\mu_2^2} (1 - e^{-\mu_2} - \mu_2 e^{-\mu_2}). \quad \square
\end{aligned}$$

Lemma 4.2 $E(D^*|T_1 = t) \leq \mu_3/(\mu_2 - p\lambda).$

Proof. $E(D^*|T_1 = t) \leq E(\mu_3 T_2|T_1 = t).$

By Theorem 3.3 and (3) we have

$$E(T_2|T_1 = t) = \int_0^\infty (\mu_2 - p\lambda) s e^{-(\mu_2 - p\lambda)s} ds, \text{ so}$$

$$E(D^*|T_1 = t) \leq \mu_3 (\mu_2 - p) \int_0^\infty s e^{-(\mu_2 - p\lambda)s} ds = \mu_3/(\mu_2 - p\lambda). \quad \square$$

Theorem 4.3 For the chosen values of $\lambda, \mu_1, \mu_2, \mu_3, p$ and t we have

$$E(T_3|T_1 = t) \geq r.$$

$$\begin{aligned}
\text{Proof. } E(M_3(a_3^+)|T_1 = t) &= (1 - p)E(D|T_1 = t) - E(D^*|T_1 = t) \\
&\quad + E(M_2(t_1^-)|T_1 = t) + E(M_3(t_1^-)|T_1 = t) + 1 \\
&> (1 - p) E(D|T_1 = t) - E(D^*|T_1 = t).
\end{aligned}$$

Thus by lemmas 4.1 and 4.2 we have

$$E(M_3(a_3^+)|T_1 = t) > \frac{(1-p)(1-\epsilon)^2(\mu_2-p\lambda)\mu_1}{\mu_2^2} (1 - e^{-\mu_2} - \mu_2 e^{-\mu_2}) - \frac{\mu_3}{\mu_2 - p\lambda}. \text{ Thus}$$

$$\begin{aligned}
E(T_3|T_1 = t) &= E(M_3(a_3^+)|T_1 = t)/\mu_3 \\
&> \frac{(1-p)(1-\epsilon)^2(\mu_2-p\lambda)\mu_1}{\mu_2^2 \mu_3} (1 - e^{-\mu_2} - \mu_2 e^{-\mu_2}) - \frac{1}{\mu_2 - p\lambda}.
\end{aligned}$$

By (4) we have $E(T_3|T_1 = t) > (1 - \epsilon)^2(r + \frac{1}{\mu_2 - p\lambda}) - \frac{1}{\mu_2 - p\lambda}$, and since ϵ was

arbitrary, we have $E(T_3|T_1 = t) \geq r$. \square

Since r was arbitrary we can choose μ_1 and t so that $E(T_3|T_1 = t) > E(T_3)$. Thus the sojourn times at node 1 and 3 are dependent for customers that go through node 2.

5. Summary. We have shown in section 4 that the sojourn times T_1 and T_3 are not independent for the network considered here. This proves the Mitrani point. Independence, of course, is sufficient to Lemoine's formula (18) as occurs, for example, in tandem queues as studied by Reich (1963). The key difference between the Lemoine result and the Reich result is that Lemoine's network allows customers to overtake each other while Reich does not. This overtaking feature allows events occurring at earlier periods of time (outputs from queue 1 that occur after our man has left queue 1) to influence events at later periods of time (the sojourn time of our man at queue 3). Clearly, if the network does not allow overtaking (e.g. if the graph of the network has at most one path from node i to node j for every i, j) then one can use results such as Reich's to show that sojourn times at nodes are mutually independent and exponentially distributed random variables.

Results related to ours have previously been discussed by Burke (1969 and 1972) and the references therein for tandem queues with multiservers. The problem discussed by Burke and that discussed by us seems to have the same underlying behavior. That is, if overtaking can occur in acyclic Jackson networks then, in general, sojourn times will not be independent.

Acknowledgements

We would like to thank Prof. Ralph L. Disney who introduced us to this problem and who discussed the contents of this paper with us during its evolution.

References

- [1] Beutler, F. J. and Melamed, B. (1978), "Decomposition and Customer Streams of Feedback Networks of Queues in Equilibrium". Oper. Res. 26, 1059-1072.
- [2] Burke, P. J. (1969) "The Dependence of Sojourn Times in Tandem M/M/s Queues". Oper. Res., 17, 754-755.
- [3] Burke, P. J. (1972) "Output Processes and Tandem Queues". Proceedings of the Symposium on Computer-Communications Networks and Teletraffic, Polytechnic Institute of Brooklyn, April 4-6, 419-428.
- [4] Çinlar, E. (1975) Introduction to Stochastic Processes. Prentice Hall, Englewood Cliffs, New Jersey.
- [5] Disney, R. L., Farrell, R. L., and de Moraes, P. R. (1973) "A Characterization of M/G/1 Queues with Renewal Departure Processes". Mgmt. Sci. 19, 1222-1228.
- [6] Jackson, J. R. (1957) "Networks of Waiting Lines". Oper. Res. 5, 518-521.
- [7] Lemoine, A. (1977) "Networks of Queues - A Survey of Equilibrium Analysis". Mgmt. Sci. 24, 464-481.
- [8] Mitrani, I. (1978) "A Critical Note on a Result of Lemoine". Private Communication.
- [9] Reich, E. (1963) "Note on Tandem Queues". Ann. Math. Stat. 34, 338-341.